

A FAMILY OF HEAT FUNCTIONS AS SOLUTIONS OF INDETERMINATE MOMENT PROBLEMS

RICARDO GÓMEZ AND MARCOS LÓPEZ-GARCÍA

ABSTRACT. We construct a family of functions satisfying the heat equation and show how they can be used to generate solutions to indeterminate moment problems. The following cases are considered: log-normal, generalized Stieltjes-Wigert and q -Laguerre.

1. INTRODUCTION

For a real-valued, measurable function f defined on $[0, \infty)$, its n th moment is defined as $s_n(f) = \int_0^\infty x^n f(x) dx$, $n \in \mathbb{N} = \{0, 1, \dots\}$. Let $(s_n)_{n \geq 0}$ be a sequence of real numbers. If f is a real-valued, measurable function defined on $[0, \infty)$ with moment sequence $(s_n)_{n \geq 0}$ we say that f is a solution to the Stieltjes moment problem (related to $(s_n)_{n \geq 0}$). If the solution is unique, the moment problem is called M -determinate. Otherwise the moment problem is said to be M -indeterminate. When we replace \mathbb{N} with \mathbb{Z} we can formulate the same problem (the so-called strong Stieltjes moment problem).

In [9] Stieltjes was the first to give examples of M -indeterminate moment problems. He showed that the log-normal distribution with density on $(0, \infty)$ given as

$$d_\sigma(x) = (2\pi\sigma^2)^{-1/2} x^{-1} \exp\left(-\frac{(\log x)^2}{2\sigma^2}\right), \quad \sigma > 0,$$

together with the densities ($a \in [-1, 1]$)

$$d_\sigma(x) (1 + a \sin(2\pi\sigma^{-2} \log x)) \geq 0,$$

have all the moment sequence $(e^{n^2\sigma^2/2})_{n \geq 0}$. So, the log-normal moment problem is M -indeterminate.

In fact, for $\beta \in \mathbb{R}$ we have

$$s_n(x^\beta d_\sigma) = q^{-(n+\beta)^2/2}, \quad n \in \mathbb{Z},$$

2000 *Mathematics Subject Classification.* Primary: 44A60, Secondary 35K05.
Key words and phrases. Indeterminate moment problem; heat function.

where $q = e^{-\sigma^2}$.

The Stieltjes' example and the work in [2] gave rise to the present paper. By looking for real-valued, measurable functions h such that

$$g_{\sigma,\beta}(x) = x^\beta d_\sigma(x) \{1 + h(\sigma^{-2} \log(xq^\beta))\}$$

satisfy $s_n(g_{\sigma,\beta}) = s_n(x^\beta d_\sigma)$ for all $n \in \mathbb{Z}$, we are faced (Proposition 1) with the problem of characterizing the real-valued, measurable functions h satisfying

$$(1) \quad \int_{\mathbb{R}} \exp(-\sigma^2 x^2/2) h(x+n) dx = 0, \quad \forall n \in \mathbb{Z}.$$

In particular, if h is a 1-periodic, real-valued, measurable function, then the last equality is equivalent to

$$(2) \quad \int_0^1 \theta(x, 2^{-1}\sigma^{-2}) h(x) dx = 0,$$

where θ is the so-called theta function given by

$$\theta(x, t) = (4\pi t)^{-1/2} \sum_{n \in \mathbb{Z}} e^{-(x+n)^2/4t}, \quad (\text{see [5, page 59]}).$$

The 1-periodic, positive function θ satisfies the heat equation on \mathbb{R}_+^2

$$\frac{\partial u}{\partial t}(x, t) = \frac{\partial^2 u}{\partial x^2}(x, t).$$

Notice that if h satisfies (1) or (2) then so does ah , $a \in \mathbb{R}$. Moreover, when h is bounded below (above) there is $a \in \mathbb{R}$ such that $1 + ah \geq 0$. Hence in this case a probability density function can be obtained by a standard normalizing procedure.

It only remains for us to find some interesting 1-periodic functions h satisfying (2). By setting $h_c(x) = q^{c^2/2} \sigma^2 M_c^{-1} \theta(x+c, 2^{-1}\sigma^{-2})^{-1} - 1$, $c \in [0, 1)$, we obtain the well-known classical solution (see e.g. [2])

$$\begin{aligned} w_c(x) &= d_\sigma(x) (1 + h_c(\sigma^{-2} \log x)) \\ &= \frac{x^{c-1}}{M_c \left(q, -q^{\frac{1}{2}-c}x, -q^{\frac{1}{2}+c}/x; q \right)_\infty} \end{aligned}$$

to the log-normal moment problem, where M_c is the constant that makes $\int_0^\infty w_c(x) dx = 1$. Information about theta functions and orthogonal polynomials can be found in [1].

To get more examples, for $\alpha \geq -1$ we define the following function

$$\theta_\alpha(x, t) = \sum_{n \in \mathbb{Z}} (2\pi n)^{2(1+\alpha)} e^{-4\pi^2 n^2 t + 2\pi n i x}.$$

Clearly θ_α is a 1-periodic function in the variable x and satisfies the heat equation on \mathbb{R}_+^2 . In addition, for $\alpha \geq -1$ and $t_1, t_2 > 0$, we show that

$$\int_0^1 \theta(y, t_1) \theta_\alpha(x - y, t_2) dy = \theta_\alpha(x, t_1 + t_2).$$

Therefore the following 1-periodic, continuous function satisfies the condition (2),

$$h_{y,t,\alpha}(x) = \theta_\alpha(y - x, t) - \theta_\alpha(y, 2^{-1}\sigma^{-2} + t), \quad y \in [0, 1], \quad t > 0.$$

Furthermore, for $\alpha > -1$ we have $\int_0^1 \theta_\alpha(x, t) dx = 0$ for all $t > 0$, thus the following 1-periodic, continuous function satisfies the condition (2),

$$h_{t,\alpha}(x) = \theta_\alpha(x, t) \theta(x, 2^{-1}\sigma^{-2})^{-1}, \quad t > 0, \quad \alpha > -1.$$

In [3, 4] Christiansen also generates new measures from old ones. The similarity of his work with the one developed here comes from the quasiperiodicity of the theta function.

The paper is organized as follows. Preliminaries are given in Section 2. We define the family $\{\theta_\alpha\}_{\alpha \geq -1}$ of heat functions in Section 3, where more functions h satisfying (2) are shown. The last two sections refer to the generalized Stieltjes-Wigert and the q -Laguerre moment problems respectively. Finally, we show a non-periodic, continuous function h fulfilling the condition (1).

2. NOTATION AND PRELIMINARIES

For $(x, t) \in \mathbb{R}_+^2$, let (see [5, pages 33, 59])

$$\begin{aligned} K(x, t) &= (4\pi t)^{-1/2} e^{-x^2/4t}, \\ \theta(x, t) &= \sum_{n \in \mathbb{Z}} K(x + n, t) \\ &= \sum_{n \in \mathbb{Z}} e^{-4\pi^2 n^2 t + 2\pi n i x}. \end{aligned}$$

The positive functions K, θ satisfy the heat equation on \mathbb{R}_+^2 . Clearly θ is a 1-periodic function in the variable x . Moreover,

$$\int_{\mathbb{R}} K(x, t) dx = 1 \quad \text{and} \quad \int_0^1 \theta(x, t) dx = 1, \quad \forall t > 0.$$

For $c \in [0, 1]$, we set

$$\mathcal{M}_c := \int_0^1 \frac{\theta(x, 2^{-1}\sigma^{-2})}{\theta(x + c, 2^{-1}\sigma^{-2})} dx.$$

Throughout this paper we will write $q = e^{-\sigma^2}$, $\sigma > 0$ fixed. The density of the log-normal distribution with parameter σ^2 can be written as

$$d_\sigma(x) = \frac{1}{x} K(\log x, 2^{-1}\sigma^2).$$

For $\beta \in \mathbb{R}$, we have

$$(3) \quad x^\beta d_\sigma(x) = q^{\beta-\beta^2/2} d_\sigma(xq^\beta).$$

Therefore,

$$\begin{aligned} s_\beta(d_\sigma) &:= \int_0^\infty x^\beta d_\sigma(x) dx = q^{-\beta^2/2} \int_0^\infty d_\sigma(x) dx \\ &= q^{-\beta^2/2} \int_{\mathbb{R}} K(x, 2^{-1}\sigma^2) dx = q^{-\beta^2/2}. \end{aligned}$$

In particular, the strong Stieltjes moment sequence of $x^\beta d_\sigma$ is given by

$$s_n(x^\beta d_\sigma) = q^{-(n+\beta)^2/2}, \quad n \in \mathbb{Z}.$$

For $0 < q < 1$, $n \in \mathbb{N}$, we introduce some notation from q -calculus (see [6, page 233]):

$$(p; q)_0 := 1, \quad (p; q)_n := \prod_{k=0}^{n-1} (1 - pq^k), \quad n \geq 1 \quad (p; q)_\infty := \prod_{k=0}^{\infty} (1 - pq^k).$$

For $\beta \in \mathbb{R}$, we set

$$(4) \quad (p; q)_\beta := \frac{(p; q)_\infty}{(pq^\beta; q)_\infty}.$$

The following easily verified identities will be used:

$$(5) \quad (p; q)_n = \frac{(p; q)_\infty}{(pq^n; q)_\infty}, \quad (p; q)_{n+\beta} = (pq^n; q)_\beta (p; q)_n.$$

We use the following notation:

$$\begin{aligned} (p_1, p_2, \dots, p_k; q)_n &= (p_1; q)_n (p_2; q)_n \cdots (p_k; q)_n, \\ (p_1, p_2, \dots, p_k; q)_\infty &= (p_1; q)_\infty (p_2; q)_\infty \cdots (p_k; q)_\infty. \end{aligned}$$

For $z \in \mathbb{C}$, we consider the two q -exponential functions,

$$(6) \quad e_q(z) = \sum_{k=0}^{\infty} \frac{z^k}{(q; q)_k} = \frac{1}{(z; q)_\infty}, \quad |z| < 1,$$

$$(7) \quad E_q(z) = \sum_{k=0}^{\infty} \frac{q^{\binom{k}{2}} z^k}{(q; q)_k} = (-z; q)_\infty.$$

For $x \in \mathbb{R}$, we define

$$L_q(x) = \sum_{n \in \mathbb{Z}} q^{\frac{1}{2}n^2} x^n.$$

The value of the sum $L_q(x)$ is known by Jacobi's triple product identity

$$L_q(x) = (q, -\sqrt{q}x, -\sqrt{q}/x; q)_\infty.$$

It is easy to check the identity

$$(8) \quad \sqrt{\frac{\sigma^2}{2\pi}} q^{x^2/2} L_q(q^{-x}) = \theta(x, 2^{-1}\sigma^{-2}), \quad \forall x \in \mathbb{R}.$$

For $c \in [0, 1)$, we introduce the following constant (see [2])

$$(9) \quad M_c := \int_0^\infty \frac{x^{c-1}}{L_q(xq^{-c})} dx = \frac{\pi q^{c(c-\frac{1}{2})} (q^c, q^{1-c}; q)_\infty}{\sin(\pi c) (q; q)_\infty^2}$$

for $c > 0$, and $M_0 = \log(q^{-1})$. By the Monotone Convergence Theorem and equality (8), we have

$$(10) \quad \begin{aligned} \mathcal{M}_c &= \sum_{n \in \mathbb{Z}} \int_0^1 \frac{K(x+n, 2^{-1}\sigma^{-2})}{\theta(x+c, 2^{-1}\sigma^{-2})} dx \\ &= \int_{\mathbb{R}} \frac{q^{x^2/2}}{q^{(x+c)^2/2} L_q(q^{-(x+c)})} dx \\ &= q^{-c^2/2} \sigma^{-2} M_c. \end{aligned}$$

Proposition 1. *Let $h \in L^1(\mathbb{R}, e^{-\sigma^2(x-n)^2/2} dx)$ for all $n \in \mathbb{N}(\mathbb{Z})$. The function $g_{\sigma,\beta}(x) = x^\beta d_\sigma(x) \{1 + h(\sigma^{-2} \log(xq^\beta))\}$ has the same (strong) Stieltjes moment sequence than $x^\beta d_\sigma$ if and only if*

$$(11) \quad \int_{\mathbb{R}} K(x, 2^{-1}\sigma^{-2}) h(x+n) dx = 0 \quad \text{for every } n \in \mathbb{N}(\mathbb{Z}).$$

Proof. By using (3) and changing variables $y = -n + \sigma^{-2} \log(xq^\beta)$ we obtain

$$s_n(g_{\sigma,\beta}) = s_n(x^\beta d_\sigma) + q^{-(\beta+n)^2/2} \int_{-\infty}^\infty K(y, 2^{-1}\sigma^{-2}) h(y+n) dy$$

and the result follows. \square

In particular, if h is a 1-periodic function in $L^1((0, 1))$, then $s_n(g_{\sigma,\beta}) = s_n(x^\beta d_\sigma)$ for all $n \in \mathbb{Z}$ if and only if

$$(12) \quad \int_0^1 \theta(x, 2^{-1}\sigma^{-2}) h(x) dx = 0.$$

Remark 2. *If h satisfies (11) or (12), then so does ah , $a \in \mathbb{R}$. Moreover, when h is bounded below (above), there is $a \in \mathbb{R}$ such that $1 + ah \geq 0$.*

Definition 3. *For $\beta \in \mathbb{R}$, let \tilde{V}_β denote the set of real-valued, measurable functions f defined on $[0, \infty)$ solving the strong moment problem*

$$s_n(f) = q^{-(n+\beta)^2/2} := s_{n,\beta}, \quad n \in \mathbb{Z}.$$

Example 4. *For $\beta \in \mathbb{R}$, $x^\beta d_\sigma \in \tilde{V}_\beta$.*

Now we want to find some interesting 1-periodic functions h satisfying (12).

Example 5. *By setting $h_c(x) = \mathcal{M}_c^{-1}\theta(x+c, 2^{-1}\sigma^{-2})^{-1} - 1$, with $c \in [0, 1)$ and using (8), (10) and (12), we obtain the classical solution*

$$\begin{aligned} w_c(x) &= d_\sigma(x) \{1 + h_c(\sigma^{-2} \log x)\} \\ &= \frac{x^{c-1}}{M_c L_q(xq^{-c})} \in \tilde{V}_0. \end{aligned}$$

Example 6. *If $f \in \tilde{V}_0$, then an easy calculation shows that*

$$q^{\beta-\beta^2/2} f(q^\beta x) \in \tilde{V}_\beta, \quad \beta \in \mathbb{R}.$$

Example 7. *Let f be a 1-periodic function integrable on $(0, 1)$. Then the function*

$$h(x) = \left(f(x) - \int_0^1 f(x) dx \right) \theta(x, 2^{-1}\sigma^{-2})^{-1}$$

satisfies (12). In particular we can put $f_t(x) = \theta(x, t)$ with $t > 0$.

To get more examples, in the next section we introduce a family of functions satisfying the heat equation for which functions fulfilling the condition (12) can be defined.

3. THE FAMILIES OF HEAT FUNCTIONS $\{K_\alpha\}_\alpha, \{\theta_\alpha\}_\alpha$.

We follow the notation in [8, Chapter 9]. For $f \in L^1(\mathbb{R})$, we define its Fourier transform as

$$(\Phi f)(\xi) = \int_{-\infty}^{\infty} f(x) e^{-ix\xi} \frac{dx}{\sqrt{2\pi}}.$$

For $\alpha \geq -1$ and $t > 0$ fixed, the function $\xi^{2(1+\alpha)}e^{-\xi^2 t}$ is in $L^1(\mathbb{R})$, so we define

$$(13) \quad \begin{aligned} K_\alpha(x, t) &= \frac{1}{\sqrt{2\pi}} \Phi^{-1} \left(\xi^{2(1+\alpha)} e^{-\xi^2 t} \right) (x) \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \xi^{2(1+\alpha)} e^{-\xi^2 t} \cos(x\xi) \frac{d\xi}{\sqrt{2\pi}}. \end{aligned}$$

Then K_α is a real-valued function that satisfies the heat equation on \mathbb{R}_+^2 . We can rewrite

$$\begin{aligned} K_\alpha(x, t) &= \frac{1}{\sqrt{\pi}} t^{-1-\alpha} K(x, t) \int_{-\infty}^{\infty} \xi^{2(1+\alpha)} e^{-\left(\xi - \frac{ix}{2\sqrt{t}}\right)^2} d\xi \\ &= \frac{1}{\sqrt{\pi}} t^{-1-\alpha} K(x, t) \int_{-\infty}^{\infty} \left(\xi + \frac{ix}{2\sqrt{t}} \right)^{2(1+\alpha)} e^{-\xi^2} d\xi. \end{aligned}$$

Therefore

$$\begin{aligned} |K_\alpha(x, t)| &\leq C_\alpha t^{-1-\alpha} K(x, t) \int_{-\infty}^{\infty} \left(\xi^{2(1+\alpha)} + \frac{x^{2(1+\alpha)}}{t^{1+\alpha}} \right) e^{-\xi^2} d\xi \\ &\leq C_\alpha t^{-1-\alpha} K(x, t) \left[1 + \frac{x^{2(1+\alpha)}}{t^{1+\alpha}} \right]. \end{aligned}$$

Since $x^\lambda e^{-x} \leq C_\lambda e^{-\frac{x}{2}}$ for $x, \lambda > 0$, we have

$$\begin{aligned} \frac{x^{2(1+\alpha)}}{(4t)^{1+\alpha}} K(x, t) &= \frac{1}{\sqrt{4\pi t}} \left(\frac{x^{2(1+\alpha)}}{(4t)^{1+\alpha}} e^{-\frac{x^2}{4t}} \right) \\ &\leq \frac{C_\alpha}{\sqrt{4\pi t}} e^{-\frac{x^2}{8t}} \\ &= C_\alpha K(x, 2t). \end{aligned}$$

Whence

$$(14) \quad |K_\alpha(x, t)| \leq C_\alpha t^{-1-\alpha} (K(x, t) + K(x, 2t)),$$

and $K_\alpha(\cdot, t) \in L^1(\mathbb{R})$ for all $t > 0$.

The convolution of $f, g \in L^1(\mathbb{R})$ is given by

$$(f * g)(x) = \int_{\mathbb{R}} f(y) g(x - y) \frac{dy}{\sqrt{2\pi}}.$$

Using the fact that $\Phi(f * g) = \Phi(f) \Phi(g)$, the definition of K_α and the inversion formula, we get

$$(15) \quad \int_{\mathbb{R}} K_\alpha(y, t_1) K_\beta(x - y, t_2) dy = K_{\alpha+\beta+1}(x, t_1 + t_2),$$

whenever $\alpha, \beta \geq -1, t_1, t_2 > 0$.

Next, for $\alpha \geq -1$ we introduce the function

$$\theta_\alpha(x, t) = \sum_{n \in \mathbb{Z}} K_\alpha(x + n, t).$$

From (14) we have the estimate

$$|\theta_\alpha(x, t)| \leq C_\alpha t^{-1-\alpha} (\theta(x, t) + \theta(x, 2t)).$$

Remark 8. From (13) we have that $K_{-1} \equiv K$, $\theta_{-1} \equiv \theta$. If $\alpha \in \mathbb{N}$, then

$$K_\alpha = (-1)^{1+\alpha} \frac{\partial^{1+\alpha} K}{\partial t^{1+\alpha}}, \quad \theta_\alpha = (-1)^{1+\alpha} \frac{\partial^{1+\alpha} \theta}{\partial t^{1+\alpha}}.$$

For $\alpha > -1$, the inversion formula implies

$$\begin{aligned} \int_0^1 \theta_\alpha(x, t) dx &= \int_{-\infty}^{\infty} \Phi^{-1} \left(\xi^{2(1+\alpha)} e^{-\xi^2 t} \right) (x) \frac{dx}{\sqrt{2\pi}} \\ &= \xi^{2(1+\alpha)} e^{-\xi^2 t} \Big|_{\xi=0} = 0, \quad \forall t > 0. \end{aligned}$$

Example 9. For $\alpha > -1$, $t > 0$, the function

$$h_{t,\alpha}(x) = \theta_\alpha(x, t) \theta(x, 2^{-1} \sigma^{-2})^{-1}$$

satisfies (12).

For $\alpha, \beta \geq -1$, $t_1, t_2 > 0$, the equality (15) implies that

$$\int_0^1 \theta_\alpha(y, t_1) \theta_\beta(x - y, t_2) dy = \theta_{\alpha+\beta+1}(x, t_1 + t_2).$$

In particular, for $\alpha \geq -1$, $\beta = -1$ we obtain

$$\begin{aligned} \int_0^1 \theta(y, t_2) \theta_\alpha(x - y, t_1) dy &= \int_0^1 \theta_\alpha(y, t_1) \theta(x - y, t_2) dy \\ &= \theta_\alpha(x, t_1 + t_2). \end{aligned}$$

Example 10. For $y \in [0, 1]$, $t > 0$ the following function fulfills (12)

$$h_{y,t,\alpha}(x) = \theta_\alpha(y - x, t) - \theta_\alpha(y, 2^{-1} \sigma^{-2} + t).$$

The following proposition gives an explicit formula for θ_α .

Proposition 11. For $\alpha > -1$,

$$\theta_\alpha(x, t) = \sum_{n \in \mathbb{Z}} (2\pi n)^{2(1+\alpha)} e^{-4\pi^2 n^2 t + 2\pi n i x}.$$

Proof. From (14) we have

$$\left| \sum_{m \in \mathbb{Z}} K_\alpha(x + m, t) \right| \leq \frac{C_\alpha}{t^{1+\alpha}} (\theta(x, t) + \theta(x, 2t)).$$

Hence the series converges uniformly on compact subsets of \mathbb{R}_+^2 and therefore it is continuous. Since the series is 1-periodic in x , it admits a representation as a Fourier series,

$$\sum_{m \in \mathbb{Z}} K_\alpha(x + m, t) = \sum_{m \in \mathbb{Z}} a_m(t) e^{2\pi m i x},$$

where convergence is in $L^2([0, 1])$. Moreover,

$$\int_0^1 \sum_{m \in \mathbb{Z}} |K_\alpha(x + m, t)| dx \leq \frac{C_\alpha}{t^{1+\alpha}} \int_{-\infty}^{\infty} (K(x, t) + K(x, 2t)) dx = \frac{C_\alpha}{t^{1+\alpha}}.$$

By the Dominated Convergence Theorem we have

$$\begin{aligned} a_m(t) &= \int_0^1 \left[\sum_{n \in \mathbb{Z}} K_\alpha(x + n, t) \right] e^{-2\pi m i x} dx \\ &= \int_{-\infty}^{\infty} \sqrt{2\pi} K_\alpha(x, t) e^{-2\pi m i x} \frac{dx}{\sqrt{2\pi}} \\ &= \left(\Phi \left[\sqrt{2\pi} K_\alpha(\cdot, t) \right] \right) (2\pi m) \\ &= (2\pi m)^{2(1+\alpha)} e^{-4\pi^2 m^2 t}. \end{aligned}$$

□

The last result implies that θ_α satisfies the heat equation on \mathbb{R}_+^2 .

Example 12. If $h(x) = \sum_{n \in \mathbb{Z}} c_n e^{2\pi n i x} \in L^2([0, 1])$, then h satisfies (12) if and only if

$$\sum_{n \in \mathbb{Z}} c_n e^{-2\pi^2 n^2 / \sigma^2} = 0.$$

4. GENERALIZED STIELTJES-WIGERT

For $0 \leq p < 1$, the generalized Stieltjes-Wigert moment problem has the following weight function on $(0, \infty)$:

$$g(x; p, q) := (p, -p\sqrt{q}/x; q)_\infty d_\sigma(x).$$

When $p = 0$, the function g is the log-normal density. The next result is based on ideas in [2].

Proposition 13. For every positive function $f \in \tilde{V}_\beta$, $\beta \geq 0$, the function $(p, -p\sqrt{q}/x; q)_\infty f(x)$ has the Stieltjes moment sequence

$$s_{n, \beta, p} := (p; q)_{n+\beta} q^{-(n+\beta)^2/2}, \quad n \in \mathbb{N}.$$

Proof. Since all the functions below are positive, using (6), (7) and (4) we have

$$\begin{aligned}
\int_0^\infty x^n g(x; p, q) f(x) dx &= (p; q)_\infty \sum_{k=0}^\infty \frac{q^{k^2/2} p^k}{(q; q)_k} \int_0^\infty x^{n-k} f(x) dx \\
&= (p; q)_\infty q^{-(n+\beta)^2/2} \sum_{k=0}^\infty \frac{(pq^{n+\beta})^k}{(q; q)_k} \\
&= (p; q)_{n+\beta} q^{-(n+\beta)^2/2}, \quad n \in \mathbb{N}.
\end{aligned}$$

In fact, the last inequality holds for $n \in \mathbb{Z}$, $\beta \in \mathbb{R}$ as long as $pq^{n+\beta} < 1$. In particular, for $p = q^{1/2}$ we obtain

$$(16) \quad \int_0^\infty x^n (q^{1/2}, -q/x; q)_\infty f(x) dx = (q^{1/2}; q)_{n+\beta} q^{-(n+\beta)^2/2}$$

for $n \in \mathbb{Z}$, $\beta \in \mathbb{R}$ whenever $q^{n+\beta+1/2} < 1$. \square

Example 14. For $\beta \geq 0$, $q^{-\beta^2/2+\beta} w_c(q^\beta x) (p, -p\sqrt{q}/x; q)_\infty$ has the moment sequence $(s_{n,\beta,p})$.

More examples can be obtained if we combine (12) and the results in Section 3.

5. q -LAGUERRE

The normalized q -Laguerre polynomials $L_m^{(\alpha)}(x; q)$ (see [7]) belong to an M -indeterminate moment problem with moments

$$S_{n,\alpha} := q^{-\alpha n - \binom{n+1}{2}} (q^{\alpha+1}; q)_n, \quad n \in \mathbb{N},$$

with $0 < q < 1$, $\alpha > -1$.

Proposition 15. Let $\alpha > -1$. For every positive function $f \in \tilde{V}_{\alpha+\frac{1}{2}}$, the function $q^{(\alpha+\frac{1}{2})^2/2} (q^{\alpha+1}, -q/x; q)_\infty f(x)$ has the Stieltjes moment sequence $S_{n,\alpha}$.

Proof. Let $f \in \tilde{V}_{\alpha+\frac{1}{2}}$. It follows from (16) with $\beta = \alpha + 1/2$, (4) and (5) that the function $q^{(\alpha+\frac{1}{2})^2/2} (q^{1/2}; q)_{\alpha+\frac{1}{2}}^{-1} (q^{1/2}, -q/x; q)_\infty f(x)$ has the moment sequence

$$q^{(\alpha+\frac{1}{2})^2/2} (q^{1/2}; q)_{\alpha+\frac{1}{2}}^{-1} (q^{1/2}; q)_{n+\alpha+\frac{1}{2}} q^{-(n+\alpha+\frac{1}{2})^2/2} = S_{n,\alpha}.$$

The result follows since $(q^{1/2}; q)_{\alpha+\frac{1}{2}}^{-1} (q^{1/2}; q)_\infty = (q^{\alpha+1}; q)_\infty$. \square

Example 16. *Example 6 with $\beta = \alpha + 1/2 > -1/2$ implies that the function*

$$q^{\alpha+\frac{1}{2}} (q^{\alpha+1}, -q/x; q)_{\infty} w_c \left(q^{\alpha+\frac{1}{2}} x \right) = \frac{q^{c(\alpha+\frac{1}{2})} (q^{\alpha+1}, -q/x; q)_{\infty} x^{c-1}}{M_c(q, -q^{\alpha+1-c}x, -q^{c-\alpha}/x; q)_{\infty}}$$

with $c \in (0, 1]$, has moment sequence $S_{n,\alpha}$. In particular, when $\alpha = c-1$ and using (9), we obtain the function

$$-\frac{\sin(\pi\alpha)}{\pi} \frac{(q; q)_{\infty}}{(q^{-\alpha}; q)_{\infty}} \frac{x^{\alpha}}{(-x; q)_{\infty}}.$$

More examples can be obtained if we combine (12) and the results in Section 3.

Finally we show a non-periodic, continuous function h fulfilling condition (11).

Example 17. *For $\gamma \in \mathbb{R} \setminus 2\pi\mathbb{Q}$, consider $h(x) = (1 + k \cos(2\pi x)) \cos \gamma x$ where*

$$k = - \int_{\mathbb{R}} e^{-\sigma^2 x^2/2} \cos(\gamma x) dx \Big/ \int_{\mathbb{R}} e^{-\sigma^2 x^2/2} \cos(\gamma x) \cos(2\pi x) dx < 0.$$

So, h is not periodic at all and satisfies (11).

ACKNOWLEDGMENTS

The work of R. Gómez was partly supported by DGAPA-PAPIIT IN120605.

REFERENCES

- [1] R. Askey, Orthogonal polynomials and theta functions, in: Theta Functions Bowdoin 1987, Part 2, Brunswick, ME, 1987, in: Proc. Sympos. Pure Math., Vol. 49, Part 2, Amer. Math. Soc., Providence, RI, 1989, 299-321.
- [2] C. Berg, From discrete to absolutely continuous solutions of indeterminate moment problems, Arab. J. Math. Sci. 4, 1998, 1-18.
- [3] J. S. Christiansen, The moment problem associated with the Stieltjes-Wigert polynomials, J. Math. Anal. Appl. 277 (2003), no. 1, 218-245.
- [4] J. S. Christiansen, The moment problem associated with the q -Laguerre polynomials, Constr. Approx. 19 (2003), no. 1, 1-22.
- [5] J.R. Cannon, The One-Dimensional Heat Equation, Addison-Wesley, New York, 1984.
- [6] G. Gasper, M. Rahman, Basic Hypergeometric Series, Cambridge University Press, Cambridge, 1990.
- [7] R. Koekoek R., R. F. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue, Report 98-17, TU-Delft, 1998, <http://aw.twi.tudelft.nl/~koekoek/askey>.
- [8] W. Rudin, Real and Complex Analysis, third edition, McGraw-Hill series in higher mathematics, 1987.

- [9] T.J. Stieltjes, Recherches sur les fractions continues, Ann. Fac. Sci. Toulouse 8, 1894, 1-122; Ann. Fac. Sci. Toulouse 9, 1895, 5-47; English version contained in T.J. Stieltjes, Collected Papers, G. van Dijk (Ed.), Vol. II, Springer, Berlin, 1993.

DR. RICARDO GÓMEZ AIZA, DR. FRANCISCO MARCOS LÓPEZ GARCÍA,
INSTITUTO DE MATEMÁTICAS,
UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO,
MÉXICO, D.F., C.P. 04510.

E-mail address: flopez@matem.unam.mx

E-mail address: rgomez@matem.unam.mx