

# SELF-FOCUSING MULTIBUMP STANDING WAVES IN EXPANDING WAVEGUIDES

NILS ACKERMANN, MÓNICA CLAPP, AND FILOMENA PACELLA

ABSTRACT. Let  $M$  be a smooth  $k$ -dimensional closed submanifold of  $\mathbb{R}^N$ ,  $N \geq 2$ , and let  $\Omega_R$  be the open tubular neighborhood of radius 1 of the expanded manifold  $M_R := \{Rx : x \in M\}$ . For  $R$  sufficiently large we show the existence of positive multibump solutions to the problem

$$-\Delta u + \lambda u = f(u) \text{ in } \Omega_R, \quad u = 0 \text{ on } \partial\Omega_R.$$

The function  $f$  is superlinear and subcritical, and  $\lambda > -\lambda_1$ , where  $\lambda_1$  is the first Dirichlet eigenvalue of  $-\Delta$  in the unit ball in  $\mathbb{R}^{N-k}$ .

## 1. INTRODUCTION

Let  $M$  be a compact  $k$ -dimensional smooth submanifold of  $\mathbb{R}^N$  without boundary,  $1 \leq k \leq N - 1$ ,  $N \geq 2$ . For  $R > 0$  define

$$\Omega_R := \bigcup_{x \in M} \left\{ Rx + v : v \in (T_x M)^\perp, |v| < 1 \right\},$$

where  $T_x M$  is the tangent space of  $M$  at  $x$ , and  $(T_x M)^\perp$  is its orthogonal complement in  $\mathbb{R}^N$ . Thus, for  $R$  large enough,  $\Omega_R$  is the open tubular neighborhood of radius 1 of the expanded manifold  $M_R := \{Rx : x \in M\}$ .

We consider the problem

$$(1.1) \quad \begin{cases} -\Delta u + \lambda u = f(u) & \text{in } \Omega_R, \\ u = 0 & \text{on } \partial\Omega_R, \end{cases}$$

for  $\lambda > -\lambda_1$ , where  $\lambda_1$  is the first Dirichlet eigenvalue of  $-\Delta$  in the unit ball in  $\mathbb{R}^{N-k}$ .

Equation (1.1) appears in nonlinear optics and models standing waves in optical waveguides. The most interesting variant for applications that aspire to exploit the nonlinear properties of the material is the *self-focusing* case where  $f(u)/u \rightarrow \infty$  as  $|u| \rightarrow \infty$ . A typical example is given by  $f(u) = u^3$ , modelling Kerr's effect. For more information on the physical background see for example [13].

Geometrically, if  $k = 1$  then  $\Omega_R$  is a *tubular guide*, i.e., an optical fiber. If  $k \geq 2$  then we say that  $\Omega_R$  is a *slab*, or, more specifically, a *shell* if  $k = N - 1$ .

For expanding annular shells, i.e., tubular neighborhoods of expanding spheres, there is a considerable body of existence results for multiple solutions that are symmetric with respect to some subgroup of  $O(N)$ , cf. [4–6, 10–12, 14]. The first existence result for multiple solutions we are aware of that does not depend on symmetries of the domain was given by Dancer and Yan [8] for expanding shells that bound convex domains, in the case of  $\lambda = 0$ .

---

Research supported by CONACYT grant 129847 and PAPIIT grant IN101209 (Mexico), and by exchange funds of the Università “La Sapienza” di Roma (Italy).

Inspired by the article of Dancer and Yan we are interested in finding positive multibump solutions to (1.1) for large  $R$ , extending their result to more general domains and to the case of  $\lambda \neq 0$ .

Set

$$F(u) := \int_0^u f(s) ds \quad \text{if } u \in \mathbb{R}$$

and assume the following hypotheses on  $f$ :

**(H1)**  $f \in C^1[0, \infty) \cap C^2(0, \infty)$ .

**(H2)**  $f(0) = f'(0) = 0$ .

**(H3)** There are  $a > 0$  and  $p_1, p_2 > 1$ ,  $p_1, p_2 < (N+2)/(N-2)$  if  $N \geq 3$ , such that  $p_1 \leq p_2$  and

$$|f''(u)| \leq a(|u|^{p_1-2} + |u|^{p_2-2}) \quad \text{for all } u > 0.$$

**(H4)**  $f(u) > 0$  for  $u > 0$ .

**(H5)** There are  $b > 0$  and  $\alpha \in (1/2, 1]$  such that for  $|u|, |v|$  small

$$|f(u+v) - f(u) - f(v)| \leq b|uv|^\alpha$$

and

$$|F(u+v) - F(u) - F(v) - f(u)v - f(v)u| \leq b|uv|^{2\alpha}.$$

It is convenient to write a point in  $\mathbb{R}^N$  as  $(\xi, \eta)$ , where  $\xi \in \mathbb{R}^k$  and  $\eta \in \mathbb{R}^{N-k}$ . Moreover, we denote by

$$\mathbb{L} := \{(\xi, \eta) \in \mathbb{R}^k \times \mathbb{R}^{N-k} : |\eta| < 1\}$$

the open cylinder or slab which is, locally, the limit of  $\Omega_R$  as  $R \rightarrow \infty$ . Ground state solutions of the limit problem

$$(1.2) \quad \begin{cases} -\Delta u + \lambda u = f(u), \\ u \in H_0^1(\mathbb{L}). \end{cases}$$

will serve as building blocks for multibump solutions. Assume that the following holds:

**(H6)** Problem (1.2) has a positive solution  $U$  which is radially symmetric in  $\xi$  and  $\eta$  separately, and which is nondegenerate, in the sense that the kernel of the space of solutions to the problem

$$-\Delta u + \lambda u = f'(U)u, \quad u \in H_0^1(\mathbb{L}),$$

has dimension  $k$ .

Some examples of nonlinearities which satisfy **(H6)** are given in [7]. Note that every positive solution of (1.2) is radially symmetric in  $\eta$  and strictly decreasing in  $|\eta|$ , by [3, Theorem 1.2].

For each  $x \in M_R$  we choose a linear isometry  $A_x \in O(N)$  which maps the tangent space  $T_x M$  onto  $\mathbb{R}^k \times \{0\}$  and  $(T_x M)^\perp$  onto  $\{0\} \times \mathbb{R}^{N-k}$ , and we set

$$(1.3) \quad \mathbb{L}_x := \{x + A_x^{-1}(z) : z \in \mathbb{L}\}.$$

We consider  $U$  to be extended by 0 to all of  $\mathbb{R}^N$ , and for each  $R > 0$  we define

$$(1.4) \quad U_{x,R}(y) := U(A_x(y-x)) \quad \text{for } y \in \mathbb{R}^N.$$

Since  $U$  is radially symmetric in  $\xi$  and in  $\eta$ , the function  $U_{x,R}$  does not depend on the choice of  $A_x \in O(N)$  as long as it satisfies  $A_x(T_x M) = \mathbb{R}^k \times \{0\}$ .

Our result is the following.

**Theorem 1.1.** *Assume that **(H1)**-**(H6)** hold. Then, for each  $n \in \mathbb{N}$  there exists  $R_n > 0$  such that for every  $R \geq R_n$  there are  $n$  points  $x_{R,1}, \dots, x_{R,n} \in M_R$  and a positive solution  $u_R$  of (1.1) of the form*

$$(1.5) \quad u_R = \sum_{i=1}^n U_{x_{R,i},R} + o(1)$$

in  $H^1(\mathbb{R}^N)$  as  $R \rightarrow \infty$ . Moreover,  $|x_{R,i} - x_{R,j}| \rightarrow \infty$  as  $R \rightarrow \infty$ , if  $i \neq j$ .

In contrast to the results on expanding annular shells mentioned above, where in general the variational problem is considered in subspaces of functions that are symmetric with respect to a subgroup of  $O(N)$ , the proof of Theorem 1.1 rests on gluing rotated translates of the positive ground state solution whose existence is demanded by condition **(H6)**. As  $R \rightarrow \infty$  the possible number of bumps becomes arbitrarily large. This basic idea is the same as the one used by Dancer and Yan in [8]. Nevertheless, our proof is slightly different and simpler, and at the same time more detailed.

To prove our result we use a Lyapunov-Schmidt reduction argument and follow closely the approach of our previous paper [1]. As is typical for this method, the proof of existence of solutions with  $n$  bumps reduces to finding critical points of the natural variational functional  $J_R$  restricted to a finite dimensional manifold  $\Sigma_R$  with boundary, which can be understood as the subset of the configuration space of  $n$  points in  $M$  one obtains when subtracting a closed neighborhood of the collision set. An element of this manifold roughly marks the centers of mass of the  $n$  bumps to be glued together. The boundary of  $\Sigma_R$  consists of tuples that contain at least two centers of mass at a prescribed minimum distance. It turns out that the value of  $J_R$  decreases near the boundary of  $\Sigma_R$ , i.e. where two masses interact, so that  $J_R$  possesses an interior maximum on  $\Sigma_R$ , a global critical point.

In [1] instead we take  $k = 1$  and build solutions out of bumps that are lined up along  $M$  with alternating signs. In that setting  $J_R$  *increases* near the boundary of  $\Sigma_R$ , so that a *minimum* exists in the interior. We also consider manifolds  $M$  with boundary and define  $\Sigma_R$  to consist of those elements from  $M^n$  that have centers of mass that stay clear of each other and of the boundary of  $M$ . Since  $J_R$  also increases as one bump approaches the boundary of  $M$ , there still exists a minimum of  $J_R$  in the interior of  $\Sigma_R$ .

When the domain is an annulus, i.e. it is a tubular neighborhood of the unit sphere, there is always a positive radial solution, which is obviously different from the multibump solutions that we find and which exists for all values of the exponent  $p > 1$ . It is natural to ask whether a similar solution exists also in the case of domains which are expanding annular shells, but different from the annulus. A positive answer in this direction is given by Bartsch, Clapp, Grossi and Pacella [2]. Moreover, an interesting open question is whether similar solutions exist in tubular neighborhoods of expanding  $k$ -dimensional manifolds ( $0 < k < N$ ).

The outline of the paper is as follows: In Section 2 we compute the required estimates and in Section 3 we describe the finite dimensional reduction and prove Theorem 1.1.

## 2. PRELIMINARY ESTIMATES

We recall some inequalities from [1].

**Lemma 2.1.** *Assume that  $\mu_k > \bar{\mu} \geq 0$  for  $k = 1, 2, 3$ . Then there is  $C > 0$  such that*

$$\int_{\mathbb{R}^N} e^{-\mu_1|x-x_1|} e^{-\mu_2|x-x_2|} dx \leq C e^{-\bar{\mu}|x_1-x_2|}$$

and

$$\int_{\mathbb{R}^N} e^{-\mu_1|x-x_1|} e^{-\mu_2|x-x_2|} e^{-\mu_3|x-x_3|} dx \leq C \exp\left(-\bar{\mu} \min_{x \in \mathbb{R}^N} \sum_{k=1}^3 |x-x_k|\right)$$

for all  $x_1, x_2, x_3 \in \mathbb{R}^N$ .

*Proof.* See [1, Lemma 2.1]. ■

**Lemma 2.2.** *Let  $\alpha$  and  $b$  be the constants from condition (H5). Given  $C \geq 1$  and  $n \in \mathbb{N}$  there is a constant  $\tilde{C} = \tilde{C}(\alpha, n, C) > 0$  such that for  $u_1, u_2, \dots, u_n \in \mathbb{R}$  with  $|u_i| \leq C$  the following inequalities hold:*

$$\left| f\left(\sum_{i=1}^n u_i\right) - \sum_{i=1}^n f(u_i) \right| \leq b \sum_{i < j} |u_i u_j|^\alpha$$

and

$$\left| F\left(\sum_{i=1}^n u_i\right) - \sum_{i=1}^n F(u_i) - \sum_{i \neq j} f(u_i) u_j \right| \leq \tilde{C} \left( \sum_{i < j} |u_i u_j|^{2\alpha} + \sum_{i < j < k} |u_i u_j u_k|^{2/3} \right).$$

*Proof.* See [1, Lemma 2.2]. ■

Set  $\mathbb{B}_s^m := \{x \in \mathbb{R}^m : |x| < s\}$ . Let  $\lambda_{1,s}$  be the first Dirichlet eigenvalue of  $-\Delta$  in  $\mathbb{B}_s^{N-k}$  and  $\vartheta_{1,s}$  be the corresponding  $L^2$ -normalized positive eigenfunction. For the solution of the limit problem (1.2) the following decay estimates hold.

**Lemma 2.3.** *There are constants  $c_1, c_2 > 0$  such that*

$$c_1 |\xi|^{-\frac{k-1}{2}} e^{-\mu|\xi|} \vartheta_{1,1}(\eta) \leq |U(\xi, \eta)| \leq c_2 |\xi|^{-\frac{k-1}{2}} e^{-\mu|\xi|} \vartheta_{1,1}(\eta) \quad \text{for all } (\xi, \eta) \in \mathbb{L},$$

where  $\mu := \sqrt{\lambda + \lambda_1}$ .

*Proof.* This follows in the same way as Eq. (7) in [7, Theorem 4]. ■

Let  $\mathbb{L}_s := \{(\xi, \eta) \in \mathbb{R}^k \times \mathbb{R}^{N-k} : |\eta| < s\}$ . The bottom of the spectrum of  $-\Delta$  in  $L^2(\mathbb{L}_s)$  with Dirichlet boundary conditions coincides with  $\lambda_{1,s}$ , cf. [1, Lemma 2.5], whose proof extends to the present case.

We fix  $r > 1$  such that  $\lambda_{1,r} + \lambda > 0$ . For  $R > 0$ ,  $x \in M_R$  and  $s \in [1, r]$ , let

$$\mathbb{L}_{s,x} := \{x + A_x^{-1}(z) : z \in \mathbb{L}_s\}$$

with  $A_x$  as in (1.3). Note that the first eigenvalue of  $-\Delta$  in  $H_0^1(\Omega_R \cap \mathbb{L}_{s,x})$  satisfies  $\lambda_1(\Omega_R \cap \mathbb{L}_{s,x}) + \lambda > 0$  for large  $R$ , since  $\Omega_R \cap \mathbb{L}_{s,x}$  is an open bounded subset of  $\mathbb{L}_{r,x}$ . We write  $V_{x,s,R}$  for the unique solution to the problem

$$(2.1) \quad \begin{cases} -\Delta u + \lambda u = f(U_{x,R}) & \text{in } \Omega_R \cap \mathbb{L}_{s,x}, \\ u = 0 & \text{on } \partial(\Omega_R \cap \mathbb{L}_{s,x}), \end{cases}$$

with  $U_{x,R}$  as in (1.4). Assumption (H4) and the maximum principle, which applies by the argument in [1, Remark 2.8], yield that  $V_{x,s,R}$  is positive for  $R$  large enough. We consider  $V_{x,s,R}$  as extended by 0 to  $\mathbb{R}^N$ . When  $s = 1$  we omit it from the notation and write  $\lambda_1, \mathbb{L}_x, V_{x,R}$  instead of  $\lambda_{1,1}, \mathbb{L}_{1,x}, V_{x,1,R}$ .

The following decay estimates hold.

**Lemma 2.4.** *For each  $s \in [1, r)$  there are positive constants  $c_3, c_4$  and  $R_0$ , independent of  $x \in M_R$ , such that all quantities*

$$|U_{x,R}(y)|, \quad |\nabla U_{x,R}(y)|, \quad |V_{x,s,R}(y)|, \quad |\nabla V_{x,s,R}(y)|,$$

are bounded by  $c_3 e^{-c_4|y-x|}$  for all  $R \geq R_0$  and almost all  $y \in \mathbb{R}^N$ . Moreover,

$$|D^2 U_{x,R}(y)| \quad \text{and} \quad |D^2 V_{x,s,R}(y)|$$

are bounded uniformly in  $\mathbb{L}_x$  and  $\Omega_R \cap \mathbb{L}_{s,x}$  respectively, independently of  $R \geq R_0$ .

*Proof.* Lemma 2.3, together with standard regularity estimates, yields the estimates for  $U_{x,R}$  and its derivatives.

To prove the estimates for  $V_{x,s,R}$  we assume without loss of generality that  $x = 0$  and that  $\mathbb{R}^k \times \{0\}$  is the tangent space to  $M_R$  at 0. Then there exists  $\tilde{c}_s > 0$  such that  $\vartheta_{1,r}(\eta) \geq \tilde{c}_s$  for all  $\eta \in \mathbb{B}_s^{N-k}$ . We write  $y \in \mathbb{L}_s$  as  $(\xi, \eta)$  with  $\xi \in \mathbb{R}^k$  and  $\eta \in \mathbb{B}_s^{N-k}$ , and set

$$W(y) := e^{-\nu|\xi|} \vartheta_{1,r}(\eta)$$

where  $\nu$  is a small positive constant, independent of  $R$ , which will be fixed next. A straightforward computation gives

$$\begin{aligned} -\Delta W(y) + \lambda W(y) &= \left( \frac{(N-1)\nu}{|\xi|} - \nu^2 + \lambda_{1,r} + \lambda \right) W(y) \\ &> (\lambda_{1,r} + \lambda - \nu^2) \tilde{c}_s e^{-\nu|\xi|}. \end{aligned}$$

Since  $\lambda_{1,r} + \lambda > 0$  we have that  $\lambda_{1,r} + \lambda - \nu^2 > 0$  if  $\nu$  is small enough. On the other hand, assumption **(H3)** on  $f$  together with Lemma 2.3 yield that

$$f(U_{x,R}) \leq \tilde{c}_1 e^{-\mu p_1 |\xi|},$$

for some large enough  $\tilde{c}_1 > 0$ . By comparison with equation (2.1) we obtain that  $V_{x,s,R} \leq \tilde{c}_2 W$  with  $\tilde{c}_2 := \tilde{c}_1 \tilde{c}_s^{-1} (\lambda_{1,r} + \lambda - \nu^2)^{-1}$  and, hence, the exponential bound on  $V_{x,s,R}$ . Regularity estimates using the results in [9] yield the estimates for its derivatives. Note that the boundary of  $\Omega_R \cap \mathbb{L}_{s,x}$  is Lipschitz and satisfies an exterior ball condition, uniformly as  $R \rightarrow \infty$ . ■

**Lemma 2.5.** *If  $s \in [1, r)$  and  $p \in (0, \infty)$  then we have the following asymptotic estimates as  $R \rightarrow \infty$ , independently of  $x \in M_R$ :*

$$(2.2) \quad \int_{\mathbb{R}^N} |V_{x,s,R} - U_{x,R}|^p dy = O(R^{-\min\{p,1\}}),$$

$$(2.3) \quad \int_{\mathbb{R}^N} |\nabla V_{x,s,R} - \nabla U_{x,R}|^2 dy = O(R^{-1}),$$

$$(2.4) \quad \int_{\mathbb{R}^N} |F(V_{x,s,R}) - F(U_{x,R})| dy = O(R^{-1}),$$

$$(2.5) \quad \int_{\mathbb{R}^N} |f(V_{x,s,R}) - f(U_{x,R})|^p dy = O(R^{-\min\{p,1\}}).$$

*Proof.* Let  $x$  be a point on the manifold  $M$ . After translation and rotation we may assume that  $x = 0$  and that  $\mathbb{R}^k \times \{0\}$  is the tangent space to  $M$  at 0. Since  $M$  is compact we may find  $\delta, \rho > 0$ , independent of  $x$ , and a smooth map  $h : \mathbb{B}_\rho^k \rightarrow \mathbb{B}_\delta^{N-k}$  such that

$$M \cap (\mathbb{B}_\rho^k \times \mathbb{B}_\delta^{N-k}) = \{(\xi, h(\xi)) : \xi \in \mathbb{B}_\rho^k\}$$

whose derivatives up to the order 3 are bounded independently of  $\xi \in \mathbb{B}_\rho^k$  and  $x \in M$ . Setting  $h_R(\xi) := Rh(\xi/R)$  we have that

$$\widetilde{M}_R := M_R \cap (\mathbb{B}_{\rho R}^k \times \mathbb{B}_{\delta R}^{N-k}) = \{(\xi, h_R(\xi)) : \xi \in \mathbb{B}_{\rho R}^k\}.$$

Following the argument given in [1, Lemma 3.2] one shows that there is a constant  $C$ , independent of  $x$ , such that

$$|h_R(\xi)| \leq \frac{C|\xi|^2}{R}$$

for all  $\xi \in \mathbb{B}_{\rho R}^k$ , and

$$\{\xi\} \times \mathbb{B}_1^{N-k}(h_R(\xi)) \subset (\{\xi\} \times \mathbb{R}^{N-k}) \cap \widetilde{\Omega}_R \subset \{\xi\} \times \mathbb{B}_{1+C(1+|\xi|^2)/R^2}^{N-k}(h_R(\xi))$$

for all  $\xi \in \mathbb{B}_{\rho R-1}^k$  and  $R$  large enough, where  $\widetilde{\Omega}_R$  is the tubular neighborhood of  $\widetilde{M}_R$  of radius 1. Consider the set

$$Q_R := \mathbb{B}_{R^{1/4}}^k \times \mathbb{B}_s^{N-k} \subset \mathbb{L}_s$$

and decompose  $\mathbb{R}^N$  as the union of the sets

$$\mathbb{R}^N \setminus Q_R, \quad Q_R \cap (\Omega_R \setminus \mathbb{L}), \quad Q_R \cap (\mathbb{L} \setminus \Omega_R), \quad Q_R \cap \mathbb{L} \cap \Omega_R.$$

Note that the integrals over  $Q_R \setminus (\mathbb{L} \cup \Omega_R)$  are zero. Using Lemma 2.4 and following closely the proof of [1, Lemma 3.2], with the obvious modifications, one obtains (2.2), (2.3), (2.4), (2.5). ■

Let

$$J_R(u) := \frac{1}{2} \int_{\Omega_R} (|\nabla u|^2 + \lambda u^2) dx - \int_{\Omega_R} F(u) dx, \quad u \in H_0^1(\Omega_R),$$

$$J_{\mathbb{L}}(u) := \frac{1}{2} \int_{\mathbb{L}} (|\nabla u|^2 + \lambda u^2) dx - \int_{\mathbb{L}} F(u) dx, \quad u \in H_0^1(\mathbb{L}),$$

be the energy functionals associated to problems (1.1) and (1.2) respectively.

**Lemma 2.6.**

$$(2.6) \quad \sup_{x \in M_R} \|V_{x,R} - U_{x,R}\|_{H^1(\mathbb{R}^N)} = O(R^{-1/2}),$$

$$(2.7) \quad \sup_{x \in M_R} |J_R(V_{x,R}) - J_{\mathbb{L}}(U)| = O(R^{-1}),$$

$$(2.8) \quad \sup_{x \in M_R} \|\nabla J_R(V_{x,R})\|_{H_0^1(\Omega_R)} = O(R^{-1/2}),$$

as  $R \rightarrow \infty$ .

*Proof.* The first two asymptotic estimates follow immediately from Lemma 2.5. To prove the third one we choose  $s \in (1, r)$  and a cut-off function  $\chi \in C^\infty(\mathbb{R}^{N-k})$  with  $\chi(\eta) = 1$  if  $|\eta| \leq 1$  and  $\chi(\eta) = 0$  if  $|\eta| \geq s$ . Fix  $R$  and  $x \in M_R$ . Assuming that  $x = 0$  and that  $\mathbb{R}^k \times \{0\}$  is the tangent space to  $M_R$  at 0, we write  $v \in H_0^1(\Omega_R)$  as  $v = v_1 + v_2$  where  $v_1(\xi, \eta) := \chi(\eta)v(\xi, \eta)$ . Then  $v_1 \in H_0^1(\Omega_R \cap \mathbb{L}_{s,x})$ ,  $\text{supp}(v_2) \subset \Omega_R \setminus \mathbb{L}_x$  and there exists a constant  $c_s$ , independent of  $R$  and  $x$ , such that  $\|v_1\|_{H^1(\mathbb{R}^N)} \leq$

$c_s \|v\|_{H^1(\mathbb{R}^N)}$  for all  $v \in H_0^1(\Omega_R)$ . Applying the definition of  $V_{x,s,R}$  and Lemma 2.5 we obtain

$$\begin{aligned} |DJ_R(V_{x,R})v| &= |DJ_R(V_{x,R})v_1| \\ &\leq |DJ_R(V_{x,s,R})v_1| + |DJ_R(V_{x,R})v_1 - DJ_R(V_{x,s,R})v_1| \\ &\leq \left| \int_{\mathbb{R}^N} (f(U_{x,R}) - f(V_{x,s,R}))v_1 \right| + O(R^{-1/2}) \|v_1\|_{H^1(\mathbb{R}^N)} \\ &\leq O(R^{-1/2}) \|v\|_{H^1(\mathbb{R}^N)}, \end{aligned}$$

as claimed. ■

For  $m = 1, 2$  we consider functions  $g_m: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  (to be fixed later) satisfying

$$(2.9) \quad g_2 < g_1,$$

$$(2.10) \quad g_m(R) \rightarrow \infty \quad \text{as } R \rightarrow \infty,$$

$$(2.11) \quad g_m(R) = o(R) \quad \text{as } R \rightarrow \infty.$$

Let  $D_{m,R}$  be the set of all points  $(x_1, \dots, x_n) \in (M_R)^n$  such that there exist  $i, j \in \{1, \dots, n\}$  with  $i \neq j$  and  $|x_i - x_j| \leq g_m(R)$ , and let

$$\mathcal{U}_{m,R} := (M_R)^n \setminus D_{m,R}.$$

Then  $\mathcal{U}_{1,R}$  and  $\mathcal{U}_{2,R}$  are open subsets of  $(M_R)^n$  with  $\overline{\mathcal{U}_{1,R}} \subset \mathcal{U}_{2,R}$ . For  $X = (x_1, \dots, x_n) \in \mathcal{U}_{2,R}$  we define  $\varphi_R: \mathcal{U}_{2,R} \rightarrow H_0^1(\Omega_R)$  by

$$(2.12) \quad \varphi_R(X) := \sum_{i=1}^n V_{x_i,R}.$$

**Proposition 2.7.** *Let  $\alpha$  be as in (H5) and fix  $\gamma \in (1/2, \alpha)$ . Then*

$$\sup_{X \in \mathcal{U}_{2,R}} \|\nabla J_R(\varphi_R(X))\|_{H_0^1(\Omega_R)} = O(e^{-\gamma\mu g_2(R)}) + O(R^{-1/2})$$

as  $R \rightarrow \infty$ .

*Proof.* Fix  $X = (x_1, \dots, x_n) \in \mathcal{U}_{2,R}$ . To simplify notation we write  $U_i := U_{x_i,R}$  and  $V_i := V_{x_i,R}$ . If  $v \in H_0^1(\Omega_R)$  satisfies  $\|v\|_{H_0^1(\Omega_R)} = 1$ , using Lemmas 2.6, 2.2, 2.5, 2.3 and 2.1, in this order, we obtain

$$\begin{aligned} |DJ_R(\varphi_R(X))v| &= \left| \sum_{i=1}^n DJ_R(\varphi_R(V_i))v + \int_{\Omega_R} \left( \sum_{i=1}^n f(V_i) - f\left(\sum_{i=1}^n V_i\right) \right) v \right| \\ &\leq \sum_{i=1}^n \|\nabla J_R(V_{x_i,R})\|_{H_0^1(\Omega_R)} + \left( \int_{\Omega_R} \left| \sum_{i=1}^n f(V_i) - f\left(\sum_{i=1}^n V_i\right) \right|^2 \right)^{1/2} \\ &\leq O(R^{-1/2}) + C \sum_{i < j} \left( \int_{\Omega_R} |V_i V_j|^{2\alpha} \right)^{1/2} \\ &= O(R^{-1/2}) + C \sum_{i < j} \left( \int_{\Omega_R} |U_i U_j|^{2\alpha} \right)^{1/2} \\ &= O(R^{-1/2}) + O(e^{-\gamma\mu g_2(R)}). \end{aligned}$$

These estimates are independent of the choice of  $X$ . ■

Set

$$E := J_{\mathbb{L}}(U)$$

and define

$$g_3(R) := g_1(R)^{-\frac{k-1}{2}} e^{-\mu g_1(R)} \quad \text{for } R > 0.$$

**Proposition 2.8.** *There exists  $\beta > 0$  such that*

$$\sup_{X \in \partial \mathcal{U}_{1,R}} J_R(\varphi_R(X)) \leq nE - \beta g_3(R) + o(g_3(R)) + O(R^{-1/2})$$

as  $R \rightarrow \infty$ .

*Proof.* If  $X \in \partial \mathcal{U}_{1,R}$  then  $|x_i - x_j| \geq g_1(R)$  for all  $i, j \in \{1, \dots, n\}$  and there exist  $i_0 \neq j_0$  with  $|x_{i_0} - x_{j_0}| = g_1(R)$ . We write  $U_i := U_{x_i, R}$  and  $V_i := V_{x_i, R}$ . By Lemma 2.3 we may choose  $\varepsilon, \rho > 0$  such that  $f(U_{i_0}) > \varepsilon$  and  $U_{j_0} \geq Cg_3(R)$  in  $B_\rho(x_{i_0}) := \{y \in \mathbb{R}^N : |y - x_{i_0}| < \rho\}$  for  $R$  large enough, independently of  $X \in \partial \mathcal{U}_{1,R}$ . Hence, for some  $\beta > 0$  and large enough  $R$  we have that

$$(2.13) \quad \int_{\mathbb{R}^N} f(U_{i_0}) U_{j_0} \geq \beta g_3(R).$$

Since  $U_i$  and  $V_i$  are uniformly bounded, using Lemma 2.2, estimate (2.2) and Lemmas 2.3 and 2.1 we obtain, as in the proof of [1, Prop. 3.5],

$$(2.14) \quad \begin{aligned} & \left| \int_{\Omega_R} \left[ F(\sum_i V_i) - \sum_i F(V_i) \right] - \sum_{i \neq j} \int_{\Omega_R} f(V_i) V_j \right| \\ & \leq C \sum_{i < j} \int_{\Omega_R} |V_i V_j|^{2\alpha} + C \sum_{i < j < k} \int_{\Omega_R} |V_i V_j V_k|^{2/3} \\ & = C \sum_{i < j} \int_{\Omega_R} |U_i U_j|^{2\alpha} + C \sum_{i < j < k} \int_{\Omega_R} |U_i U_j U_k|^{2/3} + O(R^{-2/3}) \\ & = o(g_3(R)) + O(R^{-2/3}). \end{aligned}$$

Therefore, using estimates (2.7), (2.14), (2.2), (2.5) and (2.13) we conclude that

$$\begin{aligned} J_R(\varphi_R(X)) &= \sum_i J_R(V_i) + \frac{1}{2} \sum_{i \neq j} \int_{\Omega_R} (\nabla V_i \cdot \nabla V_j + \lambda V_i V_j) \\ &\quad + \int_{\Omega_R} \left[ \sum_i F(V_i) - F(\sum_i V_i) \right] \\ &= nE + \frac{1}{2} \sum_{i \neq j} \int_{\Omega_R} f(U_i) V_j - \sum_{i \neq j} \int_{\Omega_R} f(V_i) V_j \\ &\quad + o(g_3(R)) + O(R^{-1/2}) \\ &= nE - \frac{1}{2} \sum_{i \neq j} \int_{\Omega_R} f(U_i) U_j + o(g_3(R)) + O(R^{-1/2}) \\ &\leq nE - \frac{1}{2} \int_{\Omega_R} f(U_{i_0}) U_{j_0} + o(g_3(R)) + O(R^{-1/2}) \\ &\leq nE - \beta g_3(R) + o(g_3(R)) + O(R^{-1/2}), \end{aligned}$$

as claimed.  $\blacksquare$

**Proposition 2.9.** *As  $R \rightarrow \infty$ ,*

$$\sup_{X \in \mathcal{U}_{1,R}} J_R(\varphi_R(X)) \geq nE + o(g_3(R)) + O(R^{-1/2}).$$

*Proof.* We fix  $n$  distinct points  $x_1, \dots, x_n \in M$  and set  $X_R := (Rx_1, \dots, Rx_n)$ . Then  $X_R \in \mathcal{U}_{1,R}$  for  $R$  large enough because of (2.11). As in the proof of Proposition 2.8 we have that

$$J_R(\varphi_R(X)) = nE - \frac{1}{2} \sum_{i \neq j} \int_{\Omega_R} f(U_i)U_j + o(g_3(R)) + O(R^{-1/2}).$$

Let  $\delta \in (0, \min_{i \neq j} |x_i - x_j|)$  and  $\bar{\mu} \in (0, \mu)$ . Lemmas 2.3 and 2.1 yield

$$\int_{\Omega_R} f(U_i)U_j = O(e^{-\bar{\mu}\delta R}) = o(e^{-\mu g_1(R)})$$

for all  $i \neq j$ , and our claim follows. ■

### 3. FINITE DIMENSIONAL REDUCTION AND PROOF OF THEOREM 1.1

Let  $\Sigma_R := \varphi_R(\mathcal{U}_{2,R})$ . The map  $\varphi_R$  is a  $C^2$ -immersion of  $\mathcal{U}_{2,R}$  into  $H_0^1(\Omega_R)$  (cf. [1, Lemma 4.1]) but it is not injective if  $n > 1$ . Two points in  $\mathcal{U}_{2,R}$  have the same image under  $\varphi_R$  if and only if one of them is obtained from the other by a permutation of coordinates in  $M_R^n$ . Since the group of permutations acts freely on  $\mathcal{U}_{2,R}$ , the set  $\Sigma_R$  is a  $C^2$ -submanifold of  $H_0^1(\Omega_R)$ .

For  $u \in \Sigma_R$  we denote by  $P_{u,R}$  the orthogonal projection onto the normal space  $N_u \Sigma_R := (T_u \Sigma_R)^\perp$  to  $\Sigma_R$  at  $u$ . For each  $u \in H_0^1(\Omega_R)$  we consider  $D^2 J_R(u)$  as an element of  $\mathcal{L}(H_0^1(\Omega_R))$ , i.e. as the derivative of the map  $\nabla J_R : H_0^1(\Omega_R) \rightarrow H_0^1(\Omega_R)$  at the point  $u$ .

The next two lemmas are proved in the same way as the analog statements in [1].

**Lemma 3.1.** *There are  $r_0 > 0$  and  $R_1 \geq 1$  such that for  $R \geq R_1$  and for every  $u \in \Sigma_R$  there is a unique  $v_u \in u + N_u \Sigma_R$  which satisfies  $\|u - v_u\|_{H_0^1(\Omega_R)} < r_0$  and  $P_{u,R} \nabla J_R(v_u) = 0$ . Moreover,*

$$(3.1) \quad \|u - v_u\|_{H_0^1(\Omega_R)} = O(\|\nabla J_R(u)\|_{H_0^1(\Omega_R)})$$

and

$$(3.2) \quad |J_R(u) - J_R(v_u)| = O(\|\nabla J_R(u)\|_{H_0^1(\Omega_R)}^2)$$

as  $R \rightarrow \infty$  independently of  $u \in \Sigma_R$ , and the operator  $P_{u,R} D^2 J_R(v_u)|_{N_u \Sigma_R}$  is invertible in  $\mathcal{L}(N_u \Sigma_R)$ .

*Proof.* See [1, Lemma 4.3]. ■

We now fix  $r_0 > 0$  and  $R_1 \geq 1$  as in Lemma 3.1. For  $R \geq R_1$  let  $\psi_R : \mathcal{U}_{2,R} \rightarrow H_0^1(\Omega_R)$  be given by  $\psi_R(X) := v_u$ , where  $u := \varphi_R(X)$  and  $v_u$  is given by Lemma 3.1. Define  $G_R : \mathcal{U}_{2,R} \rightarrow \mathbb{R}$  by

$$G_R := J_R \circ \psi_R.$$

**Lemma 3.2.** *For  $R \geq R_1$  the map  $G_R$  is of class  $C^1$ . If  $X \in \mathcal{U}_{2,R}$  is a critical point of  $G_R$  then  $\psi_R(X)$  is a critical point of  $J_R$ .*

*Proof.* See [1, Lemma 4.4]. ■

**Proof of Theorem 1.1.** By Lemma 3.2 it suffices to show that  $G_R$  has a local maximum in  $\mathcal{U}_{2,R}$ . Propositions 2.7, 2.8 and 2.9, together with estimate (3.2), yield the following inequalities:

$$\begin{aligned}\max G_R(\partial\mathcal{U}_{1,R}) &\leq nE - \beta g_3(R) + o(g_3(R)) + O(R^{-1/2}) + O(e^{-2\gamma\mu g_2(R)}), \\ \max G_R(\overline{\mathcal{U}_{1,R}}) &\geq nE + o(g_3(R)) + O(R^{-1/2}) + O(e^{-2\gamma\mu g_2(R)}),\end{aligned}$$

with  $\beta > 0$  and  $\gamma > 1/2$ . We choose

$$g_1(R) := \frac{1}{4\mu} \log R \quad \text{and} \quad g_2(R) := \left(\frac{1}{2} + \frac{1}{4\gamma}\right) g_1(R).$$

They clearly satisfy (2.9), (2.10) and (2.11). Note that

$$R^{-1/2} = o(g_3(R)) \quad \text{and} \quad e^{-2\gamma\mu g_2(R)} = o(g_3(R)).$$

Therefore,

$$\max G_R(\overline{\mathcal{U}_{1,R}}) > \max G_R(\partial\mathcal{U}_{1,R})$$

for large  $R$ . Hence  $G_R$  has a local maximum  $X_R$  in  $\mathcal{U}_{2,R}$ . Estimate (3.1), together with Lemma 2.6, proves (1.5). Finally,  $X_R = (x_{R,1}, \dots, x_{R,n}) \in \mathcal{U}_{1,R}$  and (2.10) yield  $|x_{R,i} - x_{R,j}| \rightarrow \infty$  as  $R \rightarrow \infty$ , if  $i \neq j$ . ■

#### REFERENCES

- [1] N. ACKERMANN, M. CLAPP, AND F. PACELLA, Alternating sign multibump solutions of nonlinear elliptic equations in expanding tubular domains, 40 pages, preprint, 2011.
- [2] T. BARTSCH, M. CLAPP, M. GROSSI, AND F. PACELLA, Asymptotically radial solutions in expanding annular domains, Preprint.
- [3] H. BERESTYCKI, L.A. CAFFARELLI, AND L. NIRENBERG, Inequalities for second-order elliptic equations with applications to unbounded domains. I, *Duke Math. J.* **81** (1996), 467–494, A celebration of John F. Nash, Jr.
- [4] J. BYEON, Existence of many nonequivalent nonradial positive solutions of semilinear elliptic equations on three-dimensional annuli, *J. Differential Equations* **136** (1997), 136–165.
- [5] F. CATRINA AND Z.Q. WANG, Nonlinear elliptic equations on expanding symmetric domains, *J. Differential Equations* **156** (1999), 153–181.
- [6] C.V. COFFMAN, A nonlinear boundary value problem with many positive solutions, *J. Differential Equations* **54** (1984), 429–437.
- [7] E.N. DANCER, Real analyticity and non-degeneracy, *Math. Ann.* **325** (2003), 369–392.
- [8] E.N. DANCER AND S. YAN, Multibump solutions for an elliptic problem in expanding domains, *Comm. Partial Differential Equations* **27** (2002), 23–55.
- [9] S.J. FROMM, Potential space estimates for Green potentials in convex domains, *Proc. Amer. Math. Soc.* **119** (1993), 225–233.
- [10] M.G. LEE AND S.S. LIN, Multiplicity of positive solutions for nonlinear elliptic equations on annulus, *Chinese J. Math.* **19** (1991), 257–276.
- [11] Y.Y. LI, Existence of many positive solutions of semilinear elliptic equations on annulus, *J. Differential Equations* **83** (1990), 348–367.
- [12] S.S. LIN, Existence of many positive nonradial solutions for nonlinear elliptic equations on an annulus, *J. Differential Equations* **103** (1993), 338–349.
- [13] C. SULEM AND P.L. SULEM, The nonlinear Schrödinger equation, Applied Mathematical Sciences, vol. 139, Springer-Verlag, New York, 1999, Self-focusing and wave collapse.
- [14] T. SUZUKI, Positive solutions for semilinear elliptic equations on expanding annuli: mountain pass approach, *Funkcial. Ekvac.* **39** (1996), 143–164.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIRCUITO EXTERIOR, CIUDAD UNIVERSITARIA, 04510 MÉXICO D.F., MEXICO.

*E-mail address:* `nils@ackermath.info`

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIRCUITO EXTERIOR, CIUDAD UNIVERSITARIA, 04510 MÉXICO D.F., MEXICO.

*E-mail address:* `mclapp@matem.unam.mx`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ “LA SAPIENZA” DI ROMA, P.LE. ALDO MORO 2, 00185 ROMA, ITALY.

*E-mail address:* `pacella@mat.uniroma1.it`